

Scotland's Rural College

## Dairy cattle in a temperate climate: the effects of weather on milk yield and composition depend on management

Hill, DL; Wall, E

*Published in:*  
Animal

*DOI:*  
[10.1017/S1751731114002456](https://doi.org/10.1017/S1751731114002456)

Print publication: 01/01/2014

*Document Version*  
Peer reviewed version

[Link to publication](#)

*Citation for pulished version (APA):*

Hill, DL., & Wall, E. (2014). Dairy cattle in a temperate climate: the effects of weather on milk yield and composition depend on management. *Animal*, 9(1), 138 - 149. <https://doi.org/10.1017/S1751731114002456>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **Dairy cattle in a temperate climate: the effects of weather on milk yield and composition depend on management**

D. L. Hill<sup>\*1</sup> and E. Wall<sup>1,2</sup>

<sup>1</sup>*Animal and Veterinary Sciences Research Group, SRUC, King's Buildings, West Mains Road, Edinburgh, EH9 3JG, UK*

<sup>2</sup>*ClimateXChange, High School Yards, Edinburgh, EH1 1LZ, UK*

\*Corresponding author: [davina.hill@sruc.ac.uk](mailto:davina.hill@sruc.ac.uk)

## **Abstract**

A better understanding of how livestock respond to weather is essential to enable farming to adapt to a changing climate. Climate change is mainly expected to impact dairy cattle through heat stress and an increase in the frequency of extreme weather events. We investigated the effects of weather on milk yield and composition (fat and protein content) in an experimental dairy herd in Scotland over 21 years. Holstein Friesian cows were either housed indoors in winter and grazed over the summer or were continuously housed. Milk yield was measured daily, resulting in 762786 test day records from 1369 individuals, and fat and protein percentage were sampled once a week, giving 89331 records from 1220 cows per trait. The relative influence of 11 weather elements, measured from local outdoor weather stations, and two indices of temperature and humidity (THI), indicators of heat stress, were compared using separate Maximum Likelihood models for each element or index. Models containing a direct measure of temperature (dry bulb, wet bulb, grass or soil temperature) or a THI provided the best fits to milk yield and fat data; wind speed and the number of hours of sunshine were most important in explaining protein content. Weather elements summarised across a week's timescale from the test day usually explained milk yield and fat content better than shorter-scale (three day, test day, test day-1) metrics. Then, examining a subset of key weather variables using REML, we found that THI, wind speed and the number of hours of sunshine influenced milk yield and composition. The shape and magnitude of these effects depended on whether animals were inside or outside on the test day. The milk yield of cows outdoors was lower at the extremes of THI than at average values, and the highest yields were obtained when THI, recorded at 0900 h, was ~55 units. Cows indoors decreased milk yield as THI increased. Fat content was lower at higher THIs

than at intermediate THIs in both environments. Protein content decreased as THI increased in animals kept indoors and outdoors, and the rate of decrease was greater when animals were outside than when they were inside. Moderate wind speeds appeared to alleviate heat stress. These results show that milk yield and composition are impacted at the upper extreme of THI under conditions currently experienced in Scotland, where animals have so far experienced little pressure to adapt to heat stress.

### **Keywords**

climate change, fat percentage, heat stress, protein percentage, THI

### **Implications**

Climate change is expected to bring about drier, hotter summers and an increased frequency of extreme weather events across Europe. Here we show that milk yield and quality decline at the upper extremes of temperature and humidity even under conditions currently experienced in Scotland. We identify the values of temperature and humidity, and of other weather elements, at which performance begins to decrease. These estimates could be used in conjunction with climate projections to help policy makers understand the likely economic impact of climate change on dairy productivity.

## Introduction

Climate change will have direct effects on livestock performance and welfare, mainly through increases in temperature and the frequency of extreme weather events, and will also affect animals indirectly through changes in the availability of fodder and pasture and the distribution of pests and parasites (Gauly *et al.*, 2013). High temperatures are associated with a greater incidence of heat stress in livestock, which can have negative effects on milk yield (Bohmanova *et al.*, 2007, Hammami *et al.*, 2013), fertility (Hansen, 2009) and health (Sanker *et al.*, 2013), and increase the risk of mortality (Vitali *et al.*, 2009). Heat stress occurs when animals experience conditions above their thermal comfort zone and are unable to dissipate enough heat to maintain thermal balance (Kadzere *et al.*, 2002). This is already costly to the dairy industry in terms of management interventions and lost productivity (St-Pierre *et al.*, 2003).

An animal's tolerance to high air temperatures depends on the amount of water vapour in the air because this influences the rate of heat loss through evaporative cooling. The association between air temperature and water vapour content can be expressed as a Temperature Humidity Index (**THI**; Thom, 1959). Milk yield in Holstein dairy cows, *Bos taurus*, is traditionally said to begin declining at around 72 THI units based on work carried out in subtropical regions (Armstrong, 1994, Ravagnolo *et al.*, 2000). Thresholds of 68 (Gauly *et al.*, 2013, Renaudeau *et al.*, 2012) or even 60 units (Bruegemann *et al.*, 2012) may, however, be more characteristic of high yielding herds in temperate zones. The genetic relationship between heat tolerance and productivity is negative (Ravagnolo and Misztal, 2000),

and dairy cattle are becoming more sensitive to heat stress due to optimisation of breeding and management practices for increased performance (Kadzere *et al.*, 2002, West *et al.*, 2003). The reduction in productivity in heat stressed cows is largely a result of reduced feed intake, but high temperatures also have a direct effect on reproductive physiology and metabolism (Renaudeau *et al.*, 2012). Cattle generate metabolic heat as a by-product of milk synthesis and so higher yielding animals experience heat stress at lower THIs than lower yielders (Kadzere *et al.* 2002).

An animal's thermal tolerance is also affected by solar radiation and the velocity of ambient air (Dikmen and Hansen, 2009, Graunke *et al.*, 2011, Hammami *et al.*, 2013), while increasing precipitation is associated with declining milk production (Stull *et al.*, 2008). Weather-related stressors could potentially affect performance immediately or have a delayed impact, and yet few studies have explored the time interval between weather events occurring and impacting milk traits (St-Pierre *et al.*, 2003). Among those that have, West *et al.*, (2003) found that the effects of mean daily THI on milk yield were greatest two out of a possible three days after THI was recorded and Bouraoui *et al.* (2002) found that mean daily THIs measured 1, 2 and 3 days before the test day had a greater effect on milk yield than test day THI. These time lags might be related to the duration of digestive processes (Gauly *et al.*, 2013).

Here we used 21 years' data from a single herd at two dairy research farms on the east and west coasts of Scotland to investigate the effects of weather on milk yield and composition (fat and protein content). The study evaluates a range of weather variables collected from Meteorological Office weather stations located on the

grounds of the farms or in the close vicinity, and two THIs that are frequently used to characterise heat stress in cattle. While the effects of heat stress on dairy cows has been well-documented in tropical and sub-tropical regions (e.g. Dikmen and Hansen, 2009, West *et al.*, 2003), a growing number of studies has also reported associations between THI and milk traits in temperate regions where tolerance to heat stress is lower (Bruegemann *et al.*, 2011, Dunn *et al.*, 2014, Hammami *et al.*, 2013). Moreover, temperatures are predicted to increase over the 21<sup>st</sup> century in southern Scotland, especially in summer, with an expected mean daily maximum temperature increase of 4.3°C by the 2080s with a very slight reduction (0-5%) in humidity (Jenkins *et al.*, 2009). We therefore aimed to (1) determine the most biologically relevant way to quantify different weather elements and two THIs with respect to measurement timescale and summary statistics (mean, maximum, minimum) and to (2) test how weather currently influences milk yield and composition in cows with and without access to grazing on the test day (management group). We hypothesised that productivity would decline under extreme weather conditions, particularly at the upper extremes of THI, and that the magnitude of the effects would depend on management.

## Material and Methods

### *Subjects, maintenance and data collection*

We studied the Langhill Holstein Friesian dairy herd, consisting of approximately 200 cows, between November 1990 and July 2011. The cattle were housed at Langhill Farm, Roslin, Midlothian (55°52'1"N, 3°10'15"W), hereafter 'Farm 1', until late June

2002 and then transferred to Crichton Royal Farm, Dumfries (55°02' N, 3°34' W), 'Farm 2', a distance of 95 km. The management systems are described for Farm 1 in Veerkamp *et al.* (1994) and for Farm 2 in Pollott and Coffey (2008). Briefly, two genetic lines were created in 1976: select (**S**) and control (**C**). S cows were bred to bulls of the highest UK genetic merit for kg fat plus protein while C cows were bred to bulls that were similar to the national average for these traits. Every year, semen from 4-5 bulls that were not closely related to the cows nor known to produce calving difficulties was obtained from nationally available stock and used to serve females from the same genetic line. Females from the two lines were managed together and allocated in equal numbers to either a High Forage (**HF**) or Low Forage (**LF**) diet system. A Total Mixed Ration (**TMR**) of blended concentrates, brewers' grain and silage was offered *ad libitum* to HF cattle in the ratio 20:5:75 total dry matter (mean proportions over a full lactation) and to LF cattle in the ratio 45:5:50. All animals received concentrates in the milking parlour. Females from the same sire were assigned to the two diet groups in equal numbers.

At Farm 1, calving took place between early September and January each year. Cows were kept indoors for approximately 200 days after calving and then grazed. Those that were still indoors at the end of June were moved outside. Most grazing occurred between April and October, inclusive, depending on the availability of pasture. At Farm 2, the HF group was grazed between April and October, and otherwise maintained indoors; LF cows were continuously housed (**CH**). Calving took place all year round for both HF and LF cows, and the majority of calves were born during the winter months. Housing at both farms consisted of conventional cubicle stalls within a single building with a corrugated metal roof and no artificial ventilation.



At Farm 1, the building had walls of slatted wood and large open doors at each end; an open ridge in the roof facilitated airflow. The building at Farm 2 had open windows along the length of one side and a gated but otherwise open section (~3m wide) on each of two opposite sides surrounding an indoor loafing area.

Cows were milked twice daily at Farm 1 and three times a day at Farm 2. Milk yield (kg) was measured and summed for each day. Fat and protein content were measured twice (Farm 1, Tuesday PM and Wednesday AM) or three times (Farm 2, Tuesday PM, Wednesday AM and midday) a week, and expressed as percentages averaged across the two or three milking events. Animals remained in the study for three lactations unless they were culled due to illness or infertility.

#### *Animal data*

We extracted milk records collected on days 4-305 of the cows' first three lactations for animals that were  $\geq 75\%$  Holstein Friesian (mean  $93.0 \pm 0.19\%$ ), discarding records collected between June 2002 and July 2003 when cows were acclimatising to Farm 2. This resulted in a dataset containing 762786 test day records for milk yield from 1369 individuals on 7073 days and 89331 weekly records from 1220 animals on 958 days for fat and protein content. The number of records for each animal ranged from 3-902 (mean  $557.6 \pm 10.68$ ) for milk yield and 3-129 (mean  $73.2 \pm 10.09$ ) for fat and protein content. Test day milk yield records were matched with weather data from the same day, and fat and protein records were matched with weather data measured on the Tuesday of the same week.

#### *Weather data*

Data on 11 weather elements (Table 1) were downloaded from the British Atmospheric Data Centre website (UK Meteorological Office, 2012). These consisted of point-samples recorded at 0900 h each day and 24h summaries (mean, minimum, maximum, total). For each element we extracted data from the closest weather station to Farm 1 for the period 1990-2002 and to Farm 2 for 2003-2011. Meteorological Office weather stations that measured most elements of interest were active on the grounds of Farm 1 until 1999 and Farm 2 for the duration of the experiment. An additional five stations  $\leq 14.4$ km from Farm 1 and one station 29km from Farm 2 were used for the remaining elements and to fill in missing values. Supplementary Table S1 provides the distances that each weather element was measured from the farms, and the elevation at which it was recorded. Using these data, we calculated  $THI_1$ :

**Equation 1**

$$THI_1 = (T_{db} + T_{wb}) \times 0.72 + 40.6$$

where  $T_{db}$  was dry bulb air temperature ( $^{\circ}\text{C}$ ) and  $T_{wb}$  was wet bulb temperature ( $^{\circ}\text{C}$ ), and  $THI_2$ :

**Equation 2**

$$THI_2 = (1.8 \times T_{db} + 32) - ((0.55 - 0.0055 \times RH) \times (1.8 \times T_{db} - 26))$$

where **RH** was relative humidity (%) (National Research Council, 1971).

As weather can have a delayed effect on biological processes, and the effects of weather depend on the timescale over which animals experience them (Bertocchi et al, 2014, Renaudeau *et al.*, 2012, West *et al.*, 2003), we explored the relationship between milk traits and all weather variables on the day the cow was milked ('test day' or **TD**), the preceding day (**TD-1**), and for the number of hours of sunshine, which was measured 0000-2359h, two days before milking (**TD-2**). We calculated a

‘moving’ mean for each daily (0900 h) point-sample over the three and seven days prior to (and including) the TD, and a moving minimum and maximum for the three variables for which 24h summaries were available (precipitation,  $T_{db}$  and sunshine). We also noted the presence versus absence of lying snow on the TD and TD-1. These methods allowed us to compare different ways of expressing the weather elements, hereafter ‘weather metrics’.

### *Statistical analysis*

Weather at Farms 1 and 2 was compared using separate Generalized Least Squares models for each weather element or index fitted by Restricted Maximum Likelihood (**REML**) from the nlme package in R version 3.0.2. (R Development Core Team, 2013). Harmonic regression allowed us to account for seasonal fluctuations in weather and we applied a first order autocorrelation structure to deal with non-independence of weather values between days.

We used Akaike’s Information Criterion (**AIC**) to determine the most biologically relevant way to express each weather element and compare the explanatory power of each element with respect to milk yield, fat content and protein content (models listed in Supplementary Table S2). AIC has been used previously to compare temperature indices in explaining milk traits (Bruegemann *et al.*, 2012, Hammami *et al.*, 2013). As the metrics for summarising a given element were closely correlated, and high proportions of shared variance can lead to unreliable estimates, we fitted each metric in a separate Linear Mixed effects Model (**LMM**) (Equation 3) using Maximum Likelihood to produce a series of non-nested models. Information Theory is an appropriate method for comparing non-nested models provided that models are

fitted to identical datasets (e.g. there are no missing values) (Burnham and Anderson, 2002). As the full dataset contained missing values where data were unavailable for the closest weather stations to a farm, we created a reduced dataset of 659918 records (86.5% of the total) and 1357 animals (99.1%) for milk yield, and 77178 records (86.4% of the total) and 1212 animals (99.3%) for fat and protein content by excluding all records with missing weather values. This dataset was used only to compare weather metrics. We fitted the following model:

**Equation 3**

$$y \sim \mu + w + \text{feed group} + \text{genetic group} + (\text{feed group} \times \text{genetic group}) + \text{management} \\ + \text{farm} + \text{lact no.} + \text{DIM} + \text{animal id} + \text{TD} + \text{ordinal calving date} + \epsilon$$

where  $y$  was the response variable (milk yield, fat or protein content, all normally distributed),  $\mu$  was the overall mean and  $w$  was a single weather metric or weather metric plus weather metric  $\times$  management interaction term; ‘feed group’ (HF or LF), ‘genetic group’ (S or C), ‘management’ on the TD (grazing or housed) and ‘farm’ (1 or 2) were two-level fixed factors, ‘lactation number’ (1, 2 or 3) was a three-level ordered factor, linear and quadratic terms of ‘**DIM**’, (Days 4-305 In Milk where day 0 was the day of calving) were covariates, animal identity, ordinal calving date and TD (continuous date from the beginning of the experiment, 1-7578) were random factors (random intercepts only) and  $\epsilon$  was the error structure. We considered farm identity to control for potential changes in management and other conditions between the two farms, and ordinal calving date (1-366) to control for the time of year that cows calved. Fitting TD as a random factor allowed us to account for temporal autocorrelation, as well as potential trends related to climate and genetic improvements over the study period. To test the hypothesis that productivity declines

in extreme weather conditions, we fitted linear, quadratic and cubic terms for all continuous weather variables (except for snow depth, precipitation and visibility which were expected to have a linear effect on milk traits), retaining lower order terms where higher order terms were significant. All continuous terms were mean-centred to reduce collinearity between polynomial terms of a given variable and to improve the interpretability of the results. LMMs were fitted using the lme4 package (Bates *et al.*, 2013) in R. We selected the ‘best’ model for each weather element based on the lowest AIC, and considered 7 AIC units to be a meaningful difference between models (Burnham *et al.*, 2011). The highest ranked model for each weather element or index was refitted using REML on the same dataset to obtain less biased parameter estimates, which were calculated using lmerTest (Kuznetsova *et al.*, 2014).

Next, we tested whether the effects of weather on milk yield and composition depended on the prevailing management type (indoors or outdoors) in a single LMM for each response variable (Equation 3) using REML. To avoid fitting variables with shared variation in the same model, weather variables were limited to precipitation, WS, sunshine, and THI<sub>2</sub>, based upon Exploratory Factor Analysis (psych package; Revelle, 2013), correlation coefficients ( $\leq 0.33$  based on TD values) and AIC rankings (see Results). For each of the three weather elements and THI, the metric belonging to the highest ranked model was used. We tested for linear effects of precipitation, and linear, quadratic, cubic and quartic effects of THI<sub>2</sub>, WS and sunshine. Non-significant interactions were removed from the models (higher order terms before lower order terms) followed by non-significant main effects using backward elimination. For each significant interaction between weather and management, a

further LMM using REML was undertaken to examine the effect size and shape of the relationship for the two management groups separately. We used differentiation to calculate the ‘turning points’ where performance began to decline for polynomial relationships between weather and milk traits based on the regression equations of the post-hoc LMMs. For models fitted by REML, we present estimates of the model coefficients ( $\beta$ ) with standard errors, t-values and *P*-values assuming significance at  $P < 0.05$ . All statistical tests are two-tailed.

## Results

### *Weather conditions at the research farms*

The UK has a maritime temperate climate with mild summers and winters. Descriptive statistics for weather at the two research farms are given in Table 1. THI<sub>1</sub> and THI<sub>2</sub> showed a strong linear correlation ( $r_p = 0.986$ ,  $t_{6873} = 495.5$ ,  $P < 0.001$ ), although THI<sub>1</sub> was higher than THI<sub>2</sub> ( $t_{6874} = 150.2$ ,  $P < 0.001$ , paired test). THI<sub>1</sub> at 0900 h was  $>60$  units across the two farms on 1114 days over the study period (16.2% of TDs), and  $>70$  units on 10 days (0.1%), and THI<sub>2</sub> at 0900 h was  $>60$  units on 626 days (9.1% of TDs) and  $>70$  units on 8 days (0.1%). THI values peaked in July and were lowest between December and February, while the number of hours of sunshine was greatest in May and lowest in December and January. The research farms received  $<1$  h sunshine over 24 h on 2343 days (33.4%) and  $>9$  h on 668 days (9.5%), and WS was  $<5$  knots at 0900 h on 2464 days (36.1%) and  $>20$  knots on 415 days (6.1%). Higher values of ppt,  $T_{db}$ ,  $T_{wb}$ , THI<sub>1</sub>, THI<sub>2</sub>,  $T_s$  and  $T_g$  were recorded at Farm 2 than at Farm 1, whereas WS, visibility, snow depth and RH were greater

at Farm 1 (Table 1). There was no difference in  $P_{MSL}$  or the number of hours of sunshine at the two farms. THI increased over the 12 years of study at Farm 1 (THI<sub>1</sub>:  $\beta = 0.17 \pm 0.04$ ,  $t = 4.34$ ,  $P < 0.001$ ; THI<sub>2</sub>:  $\beta = 0.13 \pm 0.04$ ,  $t = 2.95$ ,  $P = 0.003$ ), but did not change over the 8 years at Farm 2 (THI<sub>1</sub>:  $\beta = -0.11 \pm 0.07$ ,  $t = 1.63$ ,  $P = 0.103$ ; THI<sub>2</sub>:  $\beta = -0.13 \pm 0.08$ ,  $t = 1.64$ ,  $P = 0.101$ ). The number of hours of sunshine increased over the study period at Farm 1 ( $\beta = 0.09 \pm 0.02$ ,  $t = 4.85$ ,  $P < 0.001$ ), but did not change over the years of the study at Farm 2 ( $\beta = -0.02 \pm 0.04$ ,  $t = 0.47$ ,  $P = 0.636$ ). WS decreased over the time at Farm 1 ( $\beta = -0.21 \pm 0.05$ ,  $t = 3.90$ ,  $P < 0.001$ ), but did not change at Farm 2 ( $\beta = 0.12 \pm 0.07$ ,  $t = 1.80$ ,  $P = 0.072$ ). Precipitation did not change over the study period at Farm 1 ( $\beta = 0.02 \pm 0.03$ ,  $t = 0.49$ ,  $P = 0.625$ ) or at Farm 2 ( $\beta = 0.10 \pm 0.06$ ,  $t = 1.55$ ,  $P = 0.122$ ). Daily maximum temperatures exceeded point samples measured at 0900 h by 3.3°C ( $t_{6919} = 120.6$ ,  $P < 0.001$ ), and daily minimum temperatures were 3.7°C cooler than point-samples ( $t_{6919} = 123.0$ ,  $P < 0.001$ ).

### *Comparing the effects of weather elements and metrics on milk yield and quality*

Models testing for the effects of  $T_s$  provided the best fits to the data for both milk yield and fat content, while WS models provided the best fit to protein content data (Table 2; Supplementary Table S3). Weather elements and indices were ranked in the same order for milk yield and fat content (albeit with ties for THI<sub>1</sub>, THI<sub>2</sub> and  $T_{db}$  for fat content), but followed a different order for protein content except at the end of the scale ( $P_{MSL}$ , ppt and snow were ranked 11<sup>th</sup>, 12<sup>th</sup> and 13<sup>th</sup> across all 3 milk traits). Models testing for direct measures of temperature ( $T_s$ , THI<sub>2</sub>,  $T_{db}$ , THI<sub>1</sub>,  $T_{wb}$  and  $T_g$ ) were ranked above all other models for milk yield and fat content, and in the top 9 of 13 elements or indices for protein content. THI<sub>2</sub> showed a better fit to the data than

THI<sub>1</sub> for milk yield, but the two THIs did not differ in explanatory power for milk fat and protein (Table 2). Among models that did not contain direct temperature variables, those testing for the number of hours of sunshine (7<sup>th</sup>) and RH (8<sup>th</sup>) were ranked highest for milk yield and fat content, and the model for the number of hours of sunshine was ranked second for protein content (Table 2).

Models testing for interactions between weather and management fitted the data better or (for the effects of WS and snow on fat content, and the effects of T<sub>db</sub>, THI<sub>1</sub>, T<sub>wb</sub> and snow on protein content) not significantly worse than models without the interaction term. In all but one case (TD T<sub>s</sub>), metrics applied over a week's timescale provided better fits for milk yield than metrics applied over shorter timescales. Similarly, weekly summaries were ranked more highly (or equally highly in the cases of RH, ppt and snow) than shorter term metrics for fat content, with the exception of WS, where TD was the best metric. TD or three-day metrics were usually most effective at explaining the effects of temperature variables on protein content, while weekly summaries usually explained the effects of other weather elements on protein content better than shorter term metrics. For T<sub>db</sub>, where data were available both as 0900 h point-samples and as 24h summaries, metrics derived from point-samples ranked more highly than those based on 24h summaries for all three milk traits. Models containing metrics with higher order polynomial effects usually explained the data better than those containing lower order polynomials for milk yield and fat content, although this was less frequently the case for milk protein (Supplementary Table S3). Although models varied in explanatory power, the best metric for each weather element or index significantly influenced all three milk traits



when tested individually using REML, with the exception of snow on protein content, for which no metric was significant (Supplementary Table S4).

#### *How does weather influence milk yield in dairy cattle?*

Milk yield was influenced by two-way interactions between management and each of the individual weather variables (weekly mean THI<sub>2</sub> at 0900 h, weekly maximum number of hours of sunshine, weekly mean WS and weekly mean ppt), the interaction between diet and genetic group, and main effects of farm identity, lactation number and DIM (Table 3) as follows. When cows were outside, milk yield increased with THI to 24.0 kg at 54.9 THI units, and then decreased as THI continued to increase (Figure 1, Table 3). When cattle were indoors, by contrast, increasing THI values were associated with an overall decrease in milk yield from a local maximum of 26.5 kg of milk at 32.8 THI units. Animals outdoors increased milk yield with WS to 24.1 kg at 9.1 knots, and then gradually decreased milk yield as WS increased (Figure 1, Table 3). Those indoors increased milk yield with increasing WS when WS was low, and showed no change in milk yield at higher WS. In animals indoors and outdoors, milk yield increased and then decreased as the number of hours of sunshine increased (Table 3). Performance began to decline at lower values of sunshine when animals were indoors (26.0 kg milk at 2.4 h sunshine) than when they were outdoors (24.5 kg milk at 12.8 h sunshine (Figure 1). Cattle experienced a decrease in milk yield with increasing ppt, and the rate of decline was greater in animals outdoors ( $\beta = -0.02 \pm 0.004$ ,  $t = 5.54$ ,  $P < 0.001$ ) than indoors ( $\beta = -0.01 \pm 0.002$ ,  $t = 4.10$ ,  $P < 0.001$ ). Individuals produced more milk indoors than outdoors, at Farm 1 than Farm 2 and in later lactations than in earlier lactations, and milk production decreased over a given lactation (Table 3; Table 4). Milk yield was

greater in S than C animals (effect of genetic group in HF animals:  $\beta = 4.64 \pm 0.31$ ,  $t = 14.74$ ,  $P < 0.001$ ; effect of genetic group in LF animals:  $\beta = 4.45 \pm 0.49$ ,  $t = 9.00$ ,  $P < 0.001$ ), and in LF than HF animals (effect of feed group in C animals:  $\beta = 1.75 \pm 0.03$ ,  $t = 51.39$ ,  $P < 0.001$ ; effect of feed group in S animals:  $\beta = 2.21 \pm 0.03$ ,  $t = 74.67$ ,  $P < 0.001$ ), and the difference in milk yield between LF and HF cattle was greater in S than in C animals.

#### *How does weather influence milk fat?*

The proportion of fat in milk was influenced by two-way interactions between management and weekly mean THI<sub>2</sub> at 0900 h, management and weekly minimum sunshine, and between diet and genetic group, and main effects of TD WS, farm identity, lactation number and DIM, but not by the maximum ppt over the last three days (Table 3). Fat content showed an overall decrease with THI for animals outdoors. For animals indoors, milk fat increased to a local maximum of 3.8% at 50.2 THI units, and then decreased with THI (Figure 1, Table 3). Animals outdoors and indoors increased fat content to 4.0% at 14.4 knots and then decreased fat content as WS increased (Figure 1, Table 3). Cattle kept indoors increased fat content as the number of hours of sunshine increased, whereas cattle outdoors gradually decreased fat content as the number of hours of sunshine increased (Figure 1, Table 3). Cows produced milk with a higher proportion of fat when outdoors than indoors (Table 3; Table 4), at Farm 1 than Farm 2, and in later lactations than in earlier lactations. Milk fat decreased during the first days of a given lactation and then increased (Table 3). Fat content was greater in S than C animals (effect of genetic group in HF animals:  $\beta = 0.09 \pm 0.03$ ,  $t = 2.77$ ,  $P = 0.006$ ); effect of genetic group in LF animals:  $\beta = 0.16 \pm 0.04$ ,  $t = 4.17$ ,  $P < 0.001$ ) and in HF than LF animals (effect of

feed group in C cows:  $\beta = -0.24 \pm 0.01$ ,  $t = 18.36$ ,  $P < 0.001$ ; effect of feed group in S cows:  $\beta = -0.24 \pm 0.01$ ,  $t = 20.19$ ,  $P < 0.001$ ), and the difference in fat content between S and C cattle was greater in LF than in HF individuals.

#### *How does weather influence milk protein?*

The proportion of protein in milk was influenced by two-way interactions between management and 3 separate weather variables (mean THI<sub>2</sub> over the last 3 days, weekly mean WS, weekly mean ppt), and main effects of the weekly maximum number of hours of sunshine, diet, genetic group, farm identity, lactation number and DIM (Table 3). Protein content decreased as THI increased in animals kept outdoors and indoors, and the rate of decrease was greater when animals were outside than when they were inside (Figure 1, Table 3). Animals outdoors gradually increased protein content as WS increased, whereas protein content was not influenced by WS when animals were indoors. Examining cattle kept indoors and outdoors separately, those indoors showed a tendency to increase protein content with increasing ppt ( $\beta = 0.002 \pm 0.001$ ,  $t = 1.80$ ,  $P = 0.072$ ), but there was no effect of ppt ( $\beta = -0.0001 \pm 0.0016$ ,  $t = 0.06$ ,  $P = 0.636$ ) on protein content when cattle were outdoors. Cattle indoors and outdoors decreased protein content as the number of hours of sunshine increased (Figure 1; Table 3). Cows produced more milk protein when housed outdoors than indoors, at Farm 1 than Farm 2 and in lactations 2 and 3 than in lactation 1 (Table 3; Table 4). Protein content decreased during the first days of a given lactation and then increased (Table 3). Protein content was greater in Select than Control animals (effect of genetic group in HF animals:  $\beta = 0.05 \pm 0.01$ ,  $t = 3.48$ ,  $P < 0.001$ ; effect of genetic group in LF animals:  $\beta = 0.10 \pm 0.02$ ,  $t = 5.79$ ,  $P < 0.001$ ) and in HF than in LF cattle (effect of feed group in C animals:  $\beta = 0.04 \pm 0.01$ ,  $t =$

7.58,  $P < 0.001$ ; effect of feed group in S animals:  $\beta = 0.06 \pm 0.01$ ,  $t = 11.80$ ,  $P < 0.001$ ), and the difference in milk protein between S and C cattle was greater in LF than in HF animals.

## Discussion

A better understanding of the response of livestock to current and future weather patterns is essential to enable farming to adapt to a changing climate (Gauly *et al.*, 2013). We investigated the effects of weather over a 21 year-period on milk yield and composition under different management systems in a dairy herd at two Scottish farms. The relative influence of 11 weather elements and two THIs, indicators of heat stress, was compared. Models containing direct measures of temperature provided the best fits to milk yield and milk fat data; the number of hours of sunshine and relative humidity were also important. Models considering wind speed explained protein content best, while those containing sunshine, humidity and temperature also performed well. The importance of direct temperature metrics in explaining productivity is consistent with a wealth of studies on the impact of heat stress in dairy cattle (Renaudeau *et al.*, 2012). Relatively few studies have assessed the impact of other weather variables on milk traits, but thermal indices that account for wind speed and solar radiation perform better than those that do not (Hammami *et al.*, 2013).

In our study, weather metrics summarised across a week's timescale from the test day usually explained milk traits (particularly yield and fat content) better than shorter

scale summaries. Previous studies found that weather measured prior to the test day (up to three days before) explained test day milk traits better than weather measured on the test day (Bertocchi et al., 2014, Bouraoui et al. 2002, West et al. 2003), which may be associated with the duration of digestive processes in ruminants (Gauly et al., 2013). The higher explanatory power of longer versus shorter timescales may also reflect the greater potential for extreme weather conditions to be captured in the analysis. The pattern was less clear for protein content, with weekly, three-day and TD scales performing similarly well. This suggests that weather has a more sustained impact on milk yield and fat content than on milk protein. Although recent studies have used summaries of the three days preceding milk sampling to describe weather conditions (e.g. Lambertz et al., 2014), our results suggest that weekly summaries may be more appropriate, at least for milk yield and fat content.

The effects of weather (THI<sub>2</sub>, sunshine, wind speed and precipitation) measured from outdoor weather stations on milk yield depended on whether cattle were indoors or outdoors on the test day. Cattle that were rotated between an indoor and outdoor environment responded according to the prevailing environment and produced more milk when they were indoors than outdoors. Similarly, grazing cows produced less fat-corrected milk than animals without access to grazing in another study (Lambertz et al., 2014). We assume that these results are largely a consequence of differences in diet: animals maintained indoors in our study received *ad libitum* TMR with some forage, while those outdoors ate mainly grass. TMR maximises metabolisable energy (ME) and nutrient uptake in high producing cows and can be obtained and digested more quickly than grass (Agnew and Yan, 2000). Accordingly, many studies show an increase in milk yield with feed intake (Agnew et

469 *al.*, 1998). Further to diet effects on relative productivity, the difference in the shapes  
470 of the productivity curves for animals inside and outside is probably due to  
471 differences in weather conditions experienced by cattle in the two environments.

472  
473 When animals were outside they produced less milk during extremes of THI than  
474 during average conditions, as predicted. Other authors have reported similar  
475 declines in milk yield at low THIs or cold temperatures (Bruegemann *et al.*, 2012,  
476 Rodriguez *et al.*, 1985). The rate of decrease in milk yield in our study was greater at  
477 higher values of THI than at lower values, consistent with the idea that endotherms  
478 are more tolerant of low than high body temperatures (Hansen, 2009). Cows that  
479 were indoors showed an overall decrease in milk yield with increasing THI  
480 (measured from an outdoor weather station). In northern Europe, temperatures  
481 inside cattle buildings are 3-5°C warmer than outdoors (Seedorf *et al.*, 1998).  
482 Therefore animals indoors will be less susceptible to cold stress but may experience  
483 higher temperatures than animals outside on the same day. Indoor temperatures are  
484 also likely to increase with stocking density, although density will be lower during the  
485 summer than the winter in systems with summer grazing. It would be interesting to  
486 measure microclimatic conditions inside the barn to determine how closely the  
487 animals' immediate environment is associated with different weather elements, and  
488 how microclimate influences performance. Another question worth exploring is  
489 whether a carryover effect of weather on performance exists for animals that were  
490 recently moved indoors. Similarly, the effects of weather on animals outside may  
491 depend on how long they have been outdoors.

Dikmen and Hansen (2009) observed a weak negative relationship between a dairy cow's rectal temperature and wind speed, which together with our results on wind speed and milk yield, suggests that moderate winds can alleviate heat stress. We observed a decline in milk production with increasing precipitation, and the decline was greater in animals outdoors than indoors. Stull *et al.* (2008) also reported a decrease in milk yield in cattle as precipitation increased. Precipitation is likely to affect an animal's thermal and energy balance due to a reduction in the insulative properties of its coat after wetting and the increased energy necessary to heat a layer of moist rather than dry air trapped within the coat. High precipitation and wind speeds can increase stress levels, thus reducing the availability of energy for milk production (Webster *et al.*, 2008). Beef cattle reduced feed intake but increased rumination during wet weather (Graunke *et al.*, 2011), which implies that productivity might also be reduced on rainy days in dairy cows via feed intake. On the whole, milk yield decreased as the number of hours of sunshine increased when cattle were indoors, perhaps in response to increased radiant heat from the roof.

Weather influenced milk composition as well as yield in our study. The proportion of fat in milk showed a sharp decrease with increasing THI in animals outdoors, and was lower at the upper extreme of THI than at low and intermediate THI values when cattle were indoors. Similar to milk yield, fat content was highest at moderate wind speeds. Most previous studies also report a decrease in the proportion of fat in milk (Bouraoui *et al.*, 2002, Hammami *et al.*, 2013, Smith *et al.*, 2013) or total milk fat (Lambertz *et al.*, 2014) under conditions of heat stress or increasing temperature, although others found no effect (Knapp and Grummer, 1991, Wheelock *et al.*, 2010). While an increase in the number of sunshine hours was associated with an increase

in milk yield in cows outdoors and a decrease in milk yield in cows indoors, the inverse was true for fat content. More concentrated milk yields can arise where milk production is reduced and fat synthesis remains constant, so one possibility is that sunshine influences milk fat simply through its effects on milk yield. This could be tested by evaluating the effects of sunshine on total milk fat.

Protein content decreased as THI increased in animals kept indoors and outdoors, and the rate of decrease was greater when animals were outside than when they were inside. A decline in milk protein with THI was reported by several other authors (e.g. Bouraoui *et al.*, 2002, Bruegemann *et al.*, 2012, Gantner *et al.*, 2011, Hammami *et al.*, 2013). Our results also agree with those of Lambertz *et al.* (2014), who reported a more marked decline in total protein yield with increasing THI in cows with access to pasture than those without. The increase in milk protein content with increasing wind speed when animals were outdoors was probably due to the action of wind in alleviating heat stress, while an increasing level of radiant heat from sunshine would have contributed to heat stress.

The points at which performance began to decline with increasing THI were lower in our study than in previous work (e.g. Gauly *et al.*, 2013, Ravagnolo *et al.*, 2000) for two reasons. First, ours were calculated from daily 0900 h point-samples from local weather stations. Temperature values at 0900 are probably a slight underestimation of the mean temperature over a 24h period. Second, animals in Scotland are probably less well adapted to heat stress and are thus likely to have lower thermal tolerances than cattle in warmer climates where most work was undertaken.



Climate change models predict that temperatures will get warmer this century, leading to an increased incidence of heat stress. The statistical estimates presented here can be used in conjunction with UK Climate Projections to model the economic costs (or benefits) of climate change to milk yield and quality over the 21<sup>st</sup> century under different emissions scenarios. Such predictions about future productivity can be an important tool for informing policy. In addition, climate change is expected to bring further changes, such as a longer growing season, wetter soils and a higher incidence of disease (Gauly *et al.*, 2013), and these should also be considered. Potential decreases in productivity may be offset through changes in farming practices (adaptation), such as diet, housing or selective breeding. Future studies should investigate how genetic merit influences the effects of weather on performance.

## Conclusions

Milk yield and composition were affected by extremes of THI under conditions currently experienced in Scotland, and the shape of the relationship depended on whether animals were inside or outside. Solar radiation also impacted productivity, while moderate winds helped to alleviate heat stress. Metrics summarising weather across the week preceding the test day usually explained milk traits better than shorter-term summaries. A limitation to this study is that food intake and quality can also depend on weather, and animals consumed different diets when they were indoors and outdoors. However, diet and management system are associated under typical farming practices, so this does not reduce the practical relevance of these findings.

567 **Acknowledgements**

568 Scotland's Rural College (SRUC) receives grant-in-aid from the Scottish  
569 Government. This research was funded by the Scottish Government Rural Affairs  
570 and the Environment Portfolio Strategic Research Programme 2011-2016  
571 (Environmental Change Programme and the Climate Change Centre of Expertise,  
572 ClimateXChange). The authors wish to thank the farm staff and data managers at  
573 the Dairy Cattle Research Centre for collecting and maintaining such excellent  
574 records. The authors are grateful to Prof. Mike Coffey and two anonymous referees  
575 for helpful comments on the manuscript.

## Reference List

- Agnew RE, Yan T and Gordon FJ 1998. Nutrition of the high genetic merit dairy cow-energy metabolism studies. In *Recent Advances in Animal Nutrition* (eds PC Garnsworthy & J Wiseman), pp. 181-208. Nottingham University Press, Nottingham, UK.
- Agnew RE and Yan T 2000. Impact of recent research on energy feeding systems for dairy cattle. *Livestock Production Science* 66, 197-215.
- Armstrong DV 1994. Heat Stress Interaction with Shade and Cooling. *Journal of Dairy Science* 77, 2044-2050.
- Bates D, Maechler M, Bolker B and Walker, S 2013. lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-2. <http://lme4.r-forge.r-project.org/>
- Bertocchi L, Vitali A, Lacetera N, Nardone A, Varisco G and Bernabucci U 2014. Seasonal variations in the composition of Holstein cow's milk and temperature-humidity index relationship. *Animal* 8, 667-674.
- Bohmanova J, Misztal I and Cole J 2007. Temperature-humidity indices as indicators of milk production losses due to heat stress. *Journal of Dairy Science* 90, 1947-1956.
- Bouraoui R, Lahmar M, Majdoub A, Djemali M and Belyea R 2002. The relationship of temperature-humidity index with milk production of dairy cows in a Mediterranean climate. *Animal Research* 51, 479-491.
- Bruegemann K, Gernand E, Koenig von Borstel U and Koenig S 2012. Defining and evaluating heat stress thresholds in different dairy cow production systems. *Archiv Tierzucht* 55, 13-24
- Bruegemann K, Gernand E, von Borstel U and Koenig S 2011. Genetic analyses of protein yield in dairy cows applying random regression models with time-dependent and temperature x humidity-dependent covariates. *Journal of Dairy Science* 94, 4129-4139.
- Burnham KP and Anderson DR 2002. Information and Likelihood Theory: A Basis for Model Selection and Inference. In *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* (eds KP Burnham and DR Anderson), pp. 49-97. Springer-Verlag, New York, USA.
- Burnham KP, Anderson DR and Huyvaert KP 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* 65, 23-35.
- Dikmen S and Hansen P 2009. Is the temperature-humidity index the best indicator of heat stress in lactating dairy cows in a subtropical environment? *Journal of Dairy Science* 92, 109-116.
- Dunn RJH, Mead NE, Willett KM and Parker DE 2014. Analysis of heat stress in UK dairy cattle and impact on milk yields, *Environmental Research Letters* 9, 064006.
- Gantner V, Mijic P, Kuterovac K, Solic D and Gantner R 2011. Temperature-humidity index values and their significance on the daily production of dairy cattle. *Mljekarstvo* 61, 56-63.

- 615 Gauly M, Bollwein H, Breves G, Bruegemann K, Daenicke S, Das G, Demeler J, Hansen H,  
616 Isselstein J, Koenig S, Lohoeelter M, Martinsohn M, Meyer U, Potthoff M, Sanker C,  
617 Schroeder B, Wrage N, Meibaum B, von Samson-Himmelstjerna G, Stinshoff H and  
618 Wrenzycki C 2013. Future consequences and challenges for dairy cow production systems  
619 arising from climate change in Central Europe - a review. *Animal* 7, 843-859.
- 620 Graunke KL, Schuster T and Lidfors LM 2011. Influence of weather on the behaviour of  
621 outdoor-wintered beef cattle in Scandinavia. *Livestock Science* 136, 247-255.
- 622 Hammami H, Bormann J, M'hamdi N, Montaldo H and Gengler N 2013. Evaluation of heat  
623 stress effects on production traits and somatic cell score of Holsteins in a temperate  
624 environment. *Journal of Dairy Science* 96, 1844-1855.
- 625 Hansen PJ 2009. Effects of heat stress on mammalian reproduction. *Philosophical*  
626 *Transactions of the Royal Society B-Biological Sciences* 364, 3341-3350.
- 627 Jenkins GJ, Murphy JM, Sexton DMH, Lowe JA, Jones P and Kilsby CG 2009. UK Climate  
628 Projections: Briefing report. Exeter, UK,
- 629 Kadzere CT, Murphy MR, Silanikove N and Maltz E 2002. Heat stress in lactating dairy  
630 cows: a review. *Livestock Production Science* 77, 59-91.
- 631 Knapp DM and Grummer RR 1991. Response of Lactating Dairy-Cows to Fat  
632 Supplementation During Heat-Stress. *Journal of Dairy Science* 74, 2573-2579.
- 633 Kuznetsova A, Brockhoff, PB and Christensen, RHB 2014. lmerTest: Tests for random and  
634 fixed effects for linear mixed effect models (lmer objects of lme4 package). R package  
635 version 2.0-6. <http://CRAN.R-project.org/package=lmerTest>
- 636 Lambertz C, Sanker C and Gauly M 2014. Climatic effects on milk production traits and  
637 somatic cell score in lactating Holstein-Friesian cows in different housing systems. *Journal of*  
638 *Dairy Science* 97, 319-329.
- 639 National Research Council 1971. A guide to environmental research on animals. National  
640 Academy of Science, Washington, DC, USA.
- 641 Pollott G and Coffey M 2008. The effect of genetic merit and production system on dairy cow  
642 fertility, measured using progesterone profiles and on-farm recording. *Journal of Dairy*  
643 *Science* 91, 3649-3660.
- 644 R Development Core Team 2013. R: A Language and Environment for Statistical  
645 Computing. R Foundation for Statistical Computing, Vienna, Austria.
- 646 Ravagnolo O and Misztal I 2000. Genetic component of heat stress in dairy cattle, parameter  
647 estimation. *Journal of Dairy Science* 83, 2126-2130.
- 648 Renaudeau D, Collin A, Yahav S, de Basilio V, Gourdiene J and Collier R 2012. Adaptation to  
649 hot climate and strategies to alleviate heat stress in livestock production. *Animal* 6, 707-728.
- 650 Ravagnolo O, Misztal I and Hoogenboom G 2000. Genetic component of heat stress in dairy  
651 cattle, development of heat index function. *Journal of Dairy Science* 83, 2120-2125.

- 652 Revelle W 2013. Revelle, W. (2013) psych: Procedures for Personality and Psychological  
653 Research, Version = 1.3.10. Northwestern University, Evanston, Illinois, USA,  
654 <http://CRAN.R-project.org/package=psych>.
- 655 Rodriquez LA, Mekonnen G, Wilcox CJ, Martin FG and Krienke WA 1985. Effects of relative-  
656 humidity, maximum and minimum temperature, pregnancy, and stage of lactation on milk-  
657 composition and yield. *Journal of Dairy Science* 68, 973-978.
- 658 Sanker C, Lambertz C and Gauly M 2013. Climatic effects in Central Europe on the  
659 frequency of medical treatments of dairy cows. *Animal* 7, 316-321.
- 660 Seedorf J, Hartung J, Schroder M, Linkert KH, Pedersen S, Takai H, Johnsen JO, Metz  
661 JHM, Koerkamp PWGG, Uenk GH, Phillips VR, Holden MR, Sneath RW, Short JL, White RP  
662 and Wathes CM 1998. Temperature and moisture conditions in livestock buildings in  
663 Northern Europe. *Journal of Agricultural Engineering Research* 70, 49-57.
- 664 Smith D, Smith T, Rude B and Ward S 2013. Short communication: Comparison of the  
665 effects of heat stress on milk and component yields and somatic cell score in Holstein and  
666 Jersey cows. *Journal of Dairy Science* 96, 3028-3033.
- 667 St-Pierre NR, Cobanov B and Schnitkey G 2003. Economic losses from heat stress by US  
668 livestock industries. *Journal of Dairy Science* 86, Supplement, E52-E77.
- 669 Stull C, Messam L, Collar C, Peterson N, Castillo A, Reed B, Andersen K and VerBoort W  
670 2008. Precipitation and temperature effects on mortality and lactation parameters of dairy  
671 cattle in California. *Journal of Dairy Science* 91, 4579-4591.
- 672 Thom EC 1959. The discomfort index. *Weatherwise* 12, 57-61.
- 673 UK Meteorological Office. 2012. Met Office Integrated Data Archive System (MIDAS) Land  
674 and Marine Surface Stations Data (1853-present), NCAS British Atmospheric Data Centre.  
675 [http://badc.nerc.ac.uk/view/badc.nerc.ac.uk\\_\\_ATOM\\_\\_dataent\\_ukmo-midas](http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__dataent_ukmo-midas)
- 676 Veerkamp RF, Simm G and Oldham JD 1994. Effects of Interaction Between Genotype and  
677 Feeding System on Milk-Production, Feed-Intake, Efficiency and Body Tissue Mobilization in  
678 Dairy-Cows. *Livestock Production Science* 39, 229-241.
- 679 Vitali A, Segnalini M, Bertocchi L, Bernabucci U, Nardone A and Lacetera N 2009. Seasonal  
680 pattern of mortality and relationships between mortality and temperature-humidity index in  
681 dairy cows. *Journal of Dairy Science* 92, 3781-3790.
- 682 Webster JR, Stewart M, Rogers AR and Verkerk GA 2008. Assessment of welfare from  
683 physiological and behavioural responses of New Zealand dairy cows exposed to cold and  
684 wet conditions. *Animal Welfare* 17, 19-26.
- 685 West JW, Mullinix BG and Bernard JK 2003. Effects of hot, humid weather on milk  
686 temperature, dry matter intake, and milk yield of lactating dairy cows. *Journal of Dairy*  
687 *Science* 86, 232-242.
- 688 Wheelock JB, Rhoads RP, VanBaale MJ, Sanders SR and Baumgard LH 2010. Effects of  
689 heat stress on energetic metabolism in lactating Holstein cows. *Journal of Dairy Science* 93,  
690 644-655.

691 **Table 1** *Weather data collected by Meteorological Office stations near research farms 1 (1990 to 2002) and 2 (2003 to 2011)*

Weather			Farm 1 (4177 daily records)			Farm 2 (2896 daily records)			Farm 1 vs 2	
element/index	Recording regime	Accuracy	Mean±s.e.m	Min	Max	Mean±s.e.m	Min	Max	<i>t</i>	<i>P</i>
Precipitation ( <b>ppt</b> )	Total over 24h (0900-0900)	0.1mm	2.5±0.08	0	56.0	3.1±0.11	0	55.8	3.27	**
	PS	0.1°C	8.2±0.08	-13.0	22.4	9.7±0.10	-8.9	25.2	3.81	***
Dry bulb temperature ( <b>T<sub>db</sub></b> )	Minimum over 24h (0900-0900)	0.1°C	4.6±0.07	-14.6	17.1	6.0±0.09	-13.0	18.4	10.70	***
	Maximum over 24h (0900-0900)	0.1°C	11.5±0.08	-3.1	28.3	13.1±0.10	-4.1	30.7	9.64	***
Wet bulb temperature ( <b>T<sub>wb</sub></b> )	PS	0.1°C	6.9±0.07	-13.0	19.9	8.2±0.09	-9.3	21.3	8.95	***
THI <sub>1</sub>	See T <sub>db</sub> and T <sub>wb</sub>		51.5±0.11	21.9	70.8	53.6±0.14	27.5	73.9	9.85	***
THI <sub>2</sub>	See T <sub>db</sub> and RH		47.7±0.13	11.9	70.2	50.4±0.16	20.8	73.9	11.46	***
Grass temperature ( <b>T<sub>g</sub></b> )	Minimum over 24h (0900-0900)	0.1°C	2.5±0.08	-17.4	16.1	2.8±0.10	-16.0	17.5	2.47	*
Soil temperature	PS, 30cm below the	0.1°C	8.8±0.08	0.8	19.1	10.5±0.09	1.2	20.4	9.79	***

(T <sub>s</sub> )	surface									
Wind speed ( <b>WS</b> )	0850-0900 mean, 10m	1 knot	9.4±0.12	0	44.0	5.6±0.10	0	52.0	15.60	***
	above ground									
Visibility	PS	1m	1394.1±16.78	4	4000.0	1060.4±18.29	10.0	4000.0	8.94	***
Snow depth	PS	1cm	0.3±0.03	0	25.0	0.1±0.01	0	9.0	2.48	*
Sunshine	No. hours over 24h (0000-	0.1 h	3.5±0.05	0	15.4	3.8±0.07	0	14.7	1.83	0.068
	2359); measured using									
	Campbell-Stokes recorder									
Air pressure, mean	PS	0.1 hpa	1012.5±0.20	965.1	1047.5	1013.6±0.23	962.4	1045.1	1.05	0.294
sea level ( <b>P</b> <sub>MSL</sub> )										
Relative humidity	PS	0.1%	83.0±0.18	26.7	100	80.7±0.22	28.1	100	6.48	***
<b>(RH)</b>										

692 THI = temperature humidity index; PS = point-sample.

693 Descriptive statistics are provided for each farm, and weather between the two farms is compared using separate generalised least squares models fit by  
694 restricted maximum likelihood. Averages for THI<sub>1</sub> and THI<sub>2</sub>, which we calculated from Meteorological Office data using equations (1) and (2), respectively, are  
695 also given.

696 Recording regime indicates whether values are PSs taken at 0900 h or 24 h summaries (mean, minimum, maximum, total).

697 Two-tailed levels of statistical significance are indicated by asterisks: \*P <0.05, \*\*P <0.01 and \*\*\*P <0.001.

**Table 2** The best models for each weather element or index for milk yield, fat content and protein content based on an information-theoretic comparison of 521 Maximum Likelihood models per response variable (Supplementary Table S2 shows the full set of models compared)

Weather element	Milk yield		Fat content		Protein content	
	Rank	Unique term in best model	Rank	Unique term in best model	Rank	Unique term in best model
T <sub>s</sub>	a	TD × m	a	Weekly mean × m	e	TD × m†
THI <sub>2</sub>	b	Weekly mean × m	b	Weekly mean × m	cd	3 day mean × m†
T <sub>db</sub>	c	Weekly mean × m	b	Weekly mean × m	d	TD†
THI <sub>1</sub>	d	Weekly mean × m	b	Weekly mean × m	de	TD†
T <sub>wb</sub>	e	Weekly mean × m	c	Weekly mean × m	e	TD†
T <sub>g</sub>	f	Weekly min × m	d	Weekly min × m	c	3 day min × m
sun	g	Weekly max × m	e	Weekly min × m†	b	Weekly max × m†
RH	h	Weekly mean × m	e	TD × m†	c	Weekly mean × m†
visibility	i	Weekly mean × m	f	Weekly mean × m	g	Weekly mean × m
WS	j	Weekly mean × m	g	TD†	a	Weekly mean × m
P <sub>MSL</sub>	k	Weekly mean × m	gh	Weekly mean × m†	f	3 day mean × m†
ppt	l	Weekly max × m	hi	3 day max × m†	g	Weekly mean × m†
snow	m	Weekly mean × m	i	TD presence/absence†	h	TD-1 presence/absence†



700 All 521 models were based on Equation 3 and a single dataset of 659918 records (1357 individuals) for milk yield or 77178 records (1212  
701 individuals) for fat and protein content. Each model differed from the others in a single weather metric, the presence or absence of the weather  
702 metric  $\times$  management interaction (indicated by  $\times$  m) or order of polynomial term for the weather metric. Polynomial terms and AIC values are  
703 given in Supplementary Table S3. Models are ranked from best to worst (lowest to highest AIC) for each weather element or index (see Table 1  
704 for abbreviations); 'a' represents the highest rank, and different lower case letters indicate meaningful differences ( $\geq 7$  AIC units) in rank. †  
705 indicates that more than one model had equal support for a given weather variable; equally ranked models are listed in Supplementary Table  
706 S3. TD (test day) is the day that the cow was milked; TD-1 is the day before milking.

**Table 3** LMMs to test the effect of weather and prevailing management group (indoors or outdoors) on milk yield in 1362 Holstein Friesian cows (752674 records), fat content in 1220 cows (85134 records) and protein content in 1220 cows (87446 records) during the years 1990-2011

	Milk yield (kg)				Fat (%)				Protein (%)			
Fixed effects	$\beta$	SE	<i>t</i>	<i>P</i>	$\beta$	SE	<i>t</i>	<i>P</i>	$\beta$	SE	<i>t</i>	<i>P</i>
Intercept	24.770	0.265	93.44	***	3.919	0.030	132.13	***	3.115	0.013	243.38	***
THI <sub>2</sub>	0.042	0.006	6.80	***	-0.005	0.002	-2.85	**	-0.001	0.001	-1.56	0.120
THI <sub>2</sub> (^2)	0.015	0.001	20.48	***	-0.001	<0.001	-6.12	***	<0.001	<0.001	-0.39	0.696
THI <sub>2</sub> (^3)	<0.001	<0.001	-1.53	0.127	<0.001	<0.001	-1.90	0.058	<0.001	<0.001	-1.55	0.122
THI <sub>2</sub> (^4)	<0.001	<0.001	-9.83	***	<0.001	<0.001	2.14	*	<0.001	<0.001	-0.09	0.928
ppt	-0.008	0.003	-2.92	**	-0.001	0.001	-1.53	0.127	0.001	0.001	1.05	0.296
Sun	-0.049	0.015	-3.22	**	0.040	0.020	2.01	*	-0.007	0.001	-5.65	***
Sun (^2)	0.029	0.005	5.77	***	-0.015	0.014	-1.09	0.277	-0.001	<0.001	-2.61	**
Sun (^3)	<0.001	<0.001	1.07	0.284	0.002	0.002	1.14	0.256	<0.001	<0.001	-0.53	0.595
Sun (^4)	<0.001	<0.001	-4.13	***	<0.001	0.001	0.47	0.638	<0.001	<0.001	-0.54	0.587
WS	0.085	0.013	6.78	***	0.009	0.002	3.79	***	0.002	0.002	1.30	0.195
WS (^2)	-0.014	0.002	-8.53	***	<0.001	<0.001	0.20	0.840	<0.001	<0.001	0.02	0.985
WS (^3)	0.001	<0.001	1.46	0.146	<0.001	<0.001	-2.53	*	<0.001	<0.001	-0.15	0.881

Weather affects milk yield and composition

WS (^4)	<0.001	<0.001	0.52	0.606	<0.001	<0.001	3.30	**	<0.001	<0.001	-0.09	0.931
Diet group (LF)	1.852	0.033	55.79	***	-0.306	0.012	-25.14	***	0.052	0.004	13.84	***
Genetic group (S)	4.440	0.309	14.36	***	0.091	0.028	3.28	**	0.073	0.012	6.17	***
Farm (1)	0.774	0.119	6.49	***	0.304	0.028	11.02	***	0.093	0.013	7.22	***
Management (out)	-0.714	0.030	-23.54	***	-0.027	0.009	-2.91	**	0.009	0.004	2.27	*
Lactation number (^2)	4.985	0.016	308.06	***	0.023	0.004	5.18	***	0.033	0.002	17.04	***
Lactation number (^3)	-1.320	0.010	-126.56	***	0.005	0.003	1.72	0.086	-0.026	0.001	-19.43	***
Days in milk	-0.041	<0.001	-512.92	***	0.001	<0.001	41.74	***	0.002	<0.001	151.37	***
Days in milk (^2)	<0.001	<0.001	-89.74	***	<0.001	<0.001	66.50	***	<0.001	<0.001	63.15	***
Management x THI <sub>2</sub>	0.021	0.004	5.20	***	-0.014	0.001	-9.70	***	0.002	0.001	2.16	*
Management x THI <sub>2</sub> (^2)	-0.020	0.001	-40.32	***	<0.001	<0.001	1.21	0.228	<0.001	<0.001	0.26	0.795
Management x THI <sub>2</sub> (^3)	<0.001	<0.001	-9.68	***	<0.001	<0.001	3.04	**	<0.001	<0.001	-3.07	**
Management x THI <sub>2</sub> (^4)	<0.001	<0.001	15.92	***	<0.001	<0.001	-1.78	0.076	<0.001	<0.001	2.53	*
Management x ppt	-0.020	0.002	-13.32	***	0.001	0.001	1.60	0.110	0.003	0.001	4.04	***
Management x sun	0.249	0.009	27.21	***	-0.057	0.011	-5.39	***	0.001	0.001	0.82	0.411
Management x sun (^2)	-0.036	0.003	-11.43	***	0.027	0.007	3.89	***	<0.001	<0.001	0.07	0.947
Management x sun (^3)	-0.004	<0.001	-14.63	***	-0.003	0.001	-4.02	***	<0.001	<0.001	-0.80	0.427

Management × sun (^4)	0.001	<0.001	8.65	***	<0.001	0.001	-0.88	<i>0.377</i>	<0.001	<0.001	-1.59	<i>0.111</i>
Management × WS	0.015	0.007	2.13	*	-0.001	0.001	-1.52	<i>0.128</i>	-0.016	0.001	-15.06	***
Management × WS (^2)	-0.005	0.001	-4.91	***	<0.001	<0.001	-0.56	<i>0.577</i>	<0.001	<0.001	-0.34	<i>0.735</i>
Management × WS (^3)	0.001	<0.001	3.08	**	<0.001	<0.001	-0.14	<i>0.888</i>	<0.001	<0.001	5.10	***
Management × WS (^4)	<0.001	<0.001	-3.39	***	<0.001	<0.001	0.76	<i>0.445</i>	<0.001	<0.001	-4.62	***
Diet group × genetic group	0.557	0.039	14.11	***	0.101	0.015	6.96	***	0.011	0.006	1.74	<i>0.082</i>
<b>Random intercepts</b>	<b>%σ</b>				<b>%σ</b>				<b>%σ</b>			
Animal identity	55.4				48.2				46.3			
Ordinal calving date	7.9				1.3				4.9			
Test date	5.4				8.9				10.6			
Residual variance	31.3				41.5				38.2			

709 Linear, quadratic (^2), cubic (^3) and quartic (^4) effects were tested for where indicated. Non-significant effects that were not components of  
710 significant interactions were removed from the final models; their *P*-values are italicised. WS is wind speed and ppt is precipitation









711 **Table 4** Means  $\pm$  standard errors (s.e.m) with the numbers of records and unique individuals for milk yield and fat and protein content.

712 Significant differences between levels are indicated in Table 3

		Milk yield (kg)				Fat content (%)				Protein content (%)		
		mean	s.e.m	records	cows	mean	s.e.m	records	cows	mean	s.e.m	records
Diet group	HF	23.8	0.17	435074	1026	4.2	0.02	45592	865	3.3	0.01	46865
	LF	29.4	0.24	317600	923	3.9	0.02	39542	707	3.3	0.01	40582
Genetic group	S	29.2	0.22	412594	742	4.1	0.02	44338	654	3.3	0.01	45418
	C	24.8	0.25	340080	620	3.9	0.02	40796	566	3.2	0.01	42418
Prevailing management	in	28.8	0.18	499575	1346	4.0	0.02	58625	1192	3.2	0.01	60131
	out	22.2	0.17	253099	971	4.2	0.02	26509	836	3.3	0.01	27315
Farm	1	25.5	0.27	421620	742	4.2	0.03	40025	601	3.2	0.01	39993
	2	24.8	0.27	331054	667	3.9	0.03	45109	664	3.1	0.01	47453
Lactation no.	1	20.7	0.27	327348	1300	4.0	0.03	38503	1145	3.1	0.01	39480
	2	25.9	0.27	244721	985	4.1	0.03	27273	855	3.2	0.01	28088
	3	27.8	0.26	180605	723	4.1	0.03	19358	606	3.2	0.01	19878
Overall		27.2	0.17	752674	1362	4.0	0.02	85134	1220	3.2	0.01	87446

713 The number of animals used for analyses of protein content was the same as for analyses of fat content

714 **Figure 1** The effects of i) THI, ii) wind speed ('WS') and iii) sunshine on a) daily milk yield ( $N = 752674$  records from 1362 cows), b)  
715 milk fat ( $N = 85134$  records from 1220 cows) and c) milk protein ( $N = 87446$  records from 1220 cows) in a herd of dairy cattle on two  
716 research farms in Scotland depended on whether the animals were indoors (thin unbroken line) or outdoors (thick line), except  
717 where both groups of cattle are represented by a single broken line. Weather values were recorded from the closest outdoor  
718 weather station to each farm for each element. All plots are adjusted for the terms in Equation 3, where significant, and statistical  
719 estimates for the effects presented here are provided in Table 3. Note that plots are truncated to exclude the highest and lowest  
720 0.5% of weather records due to small samples for extreme weather events.

**a.i****a.ii****a.iii****b.i****b.ii****b.iii****c.i****c.ii****c.iii**